



## Clinico-Etiological Profile of Acute Febrile Illness in Children with Sickle Cell Anemia

Dipak Madavi<sup>1</sup>, Bhagyashree Tirpude<sup>2</sup>, Shamama Subuhi<sup>3</sup>

<sup>1</sup>Associate Professor, <sup>2,3</sup>Assistant Professor

Department of Pediatrics, Indira Gandhi Govt. Medical College, Nagpur

**\*Corresponding Author:**

**Bhagyashree Tirpude**

Department of Pediatrics, Indira Gandhi Govt. Medical College, Nagpur

Type of Publication: Original Research Paper

Conflicts of Interest: Nil

### Abstract

**Objective:** In Sickle Cell anemia, fever is a common presenting symptom and is frequently the first indication of serious and life-threatening bacterial infection. Infection is a significant contributor to morbidity and mortality in Sickle Cell anemia.

High prevalence of Sickle Cell anemia & lack of study in central India showing causes of fever in them, we studied various causes of fever in patients with Sickle Cell anemia and to provide data which will help to reduce morbidity and mortality by early intervention in patients of Sickle Cell disease.

**Methods:** This was a cross sectional observational study which contains 163 patients of Sickle Cell anemia (SS) presenting with fever in the age group between 6 months to 12 years. All cases were investigated with CBC, blood culture, urine examination, chest x-ray, Blood and Urine Culture and other investigations as indicated.

**Result:** Amongst 163 cases enrolled, most common system to involve was respiratory system (44%), followed by gastrointestinal system (24%) and genitourinary system (8%). In these Pneumonia was the most common clinical diagnosis (23%).

**Conclusion:** We concluded that, the respiratory system was the most common one to get involved in a patient of sickle cell anaemia presenting with fever and pneumonia was the most common diagnosis in it.

**Keywords:** Sickle cell anemia, Fever, Bacteremia.

### INTRODUCTION

Sickle Cell Disease is an inherited hemoglobinopathy resulting from the inheritance of HbS gene either homozygously (also known as Sickle Cell Anemia), or as a compound heterozygote with other interacting abnormal hemoglobin gene (1). Sickle cell disease (SCD) is prevalent in central Africa, Middle East and Saudi Arabia, Mediterranean and in parts of India (1). SCD has a high prevalence in India and poses considerable health burden (2,3).

Fever is a common presenting symptom amongst various manifestations of Sickle cell anemia. In particular, fever is frequently the first indication of serious and life-threatening bacterial infection.

Infection is a significant contributor to morbidity and mortality in Sickle cell anemia (4,5).

There is lack of study in central India showing causes of fever in Sickle cell disease. So this study was planned to identify various infectious causes of fever in patients with Sickle cell anemia presenting with fever and to provide data which will help to reduce rates of morbidity and mortality by early intervention in patients of Sickle cell disease.

### METHODS & MATERIAL:

This study was a cross sectional observational study. We enrolled 163 patients of Sickle cell anaemia (SS pattern) in the age group of 6 months to 12 years admitted with fever  $\geq 100^{\circ}\text{F}$  in Pediatric wards of

tertiary care hospital in central India. The study period was from December 2017 to September 2019.

Considering the prevalence of 31% of acute febrile illness in Sick cell anaemia from previous studies(R), absolute allowable error of 7%, and normal deviate of 1.96% the minimum required sample size (n) was 163. The study was approved by the institutional ethics committee and a written informed consent was obtained from parents of all participants.

At the time of enrolment detailed history, physical examination and investigations were done like complete blood count (including the haemoglobin estimation, total and differential leukocyte counts) peripheral smear, blood culture, urine culture, CXR, retic count and other investigations were done as indicated.

## RESULT

In this study, amongst 163 patients enrolled, respiratory system was the most commonly involved with 72(44.17%) cases, followed by 39(23.92%) cases of gastrointestinal system, then 13(7.97%) cases of genitourinary system and 3(1.84%) cases of central nervous system.

Table no 1 shows, frequency of various clinical diagnosis found in patients of Sick cell anaemia with fever in which the respiratory system was divided into lower respiratory tract infection(LRTI) and upper respiratory tract infection(URTI), in this LRTI was further divided into lobar pneumonia which appeared in 30 patients, and bronchopneumonia which appeared in 7 patients. URTI was also further divided into pharyngitis which appeared in 23 patients and 6 patients were presented with tonsillitis. Hyper reactive airway disease (HRAD)/asthma was the

presentation in 4 patients. 2 patients found to have pulmonary tuberculosis and were receiving AKT.

Hepatitis antibody were positive in 4 patients and 16 patients had enteric fever with positive widal test, out of this salmonella was isolated in blood culture of 3 patients. Acute gastroenteritis was present in 10 patients with positive stool microscopy while 7 patients had splenic abscess. 2 patients had peritonitis out of these 1patient died in whom streptococcus pneumoniae was isolated from peritoneal fluid.

Clinically and radiologically diagnosed osteomyelitis was found in 20 patients, in which salmonella typhi was isolated from blood culture of 2 patients in which 1 had concomitant enteric fever, and also staphalococcus was isolated from blood culture of 2 patients.

Urinary tract infection was found in 13 patients with clinical features of UTI and urine microscopy showed pus cells. Out of 13 patients, 3 were urine culture positive, E. coli was isolated in 2 patients and klebsiella was isolated from urine culture of 1 patient.

7 patients had dengue fever who were diagnosed clinically and serologically. 4 patients of malaria out of which 3 had positive for plasmodium vivax and 1 was positive for plasmodium falciparum. 3 patients were reactive for HIV.

Meningitis was the diagnosis in 2 patients in which CSF study shown increased WBC but no organism was isolated in CSF culture, and one patient presented with stroke. Mumps and Measles was the clinical diagnosis in 1 patient each and 1 patient had septicemia in which E. coli was isolated in blood culture.

**TABLE NO. 1: DISTRIBUTION OF CLINICAL DIAGNOSIS IN SICKLE CELL ANEMIA WITH FEVER**

CLINICAL DIAGNOSIS.	NO OF CASES
LRTI	37 (22.63%)
- Lobar pneumonia	30 (18.4%)
- Bronchopneumonia	7 (4.2%)
URTI	29 (17.79%)
- Pharyngitis	23 (14.11%)
-Tonsillitis	

	6 (3.68%)
HRAD	4 (2.4%)
Pulmonary tuberculosis	2 (1.2%)
Enteric Fever	16 (9.8%)
Acute Gastroenteritis	10(6.13%)
Splenic abscess	7(4.2%)
Hepatitis	4(2.4%)
- Hepatitis A	4(2.4%)
- Hepatitis B	0
Peritonitis	2(1.2%)
Osteomyelitis	20(12.2%)
UTI	13(7.9%)
Dengue fever	7(4.2%)
Malaria	4(2.4%)
- Plasmodium Vivax	3(1.8%)
- Plasmodium Falciparum	1(0.6%)
HIV	3(1.8%)
Meningitis	2(1.2%)
Measles	1(0.6%)
Mumps	1(0.6%)
Septicemia	1(0.6%)
Total	163 (100%)

Table no 2 shows the distribution of bacterial pathogens isolated from body fluids, in which amongst 163 enrolled patient's blood culture was positive in 9.8% cases, urine culture was positive in 1.8% cases and peritoneal fluid culture was positive in 0.6% cases. We found streptococcus pneumoniae in 3 patients of bacteremia, while it was also isolated from 1 peritoneal fluid culture. Klebsiella pneumoniae was positive in 3 patients of bacteremia and in 1 urine culture. E. coli was found in 2 patients of bacteremia and also isolated in 2 urine culture. Salmmonella typhi was found in 3 patients of bacteremia. Staphalococcus aurease and Coagulase negative staphylococcus were found in 2-2 patients of bacteremia. Pseudomonas aeuroginosa was found in 1 patient of bacteremia. Culture of CSF were sterile in 2 patients of meningitis.

**TABLE NO. 2: DISTRIBUTION OF DIFFERENT BACTERIAL PATHOGENS**

ORGANISM	BLOOD	URINE	PERITONEA L FLUID	CSF	TOTAL
Streptococcus pneumonia	3	0	1	0	4(2.4%)
Klebsiellapneumoniae	3	1	0	0	4(2.4%)
Salmonella typhi	3	0	0	0	3(1.8%)
E.coli	2	2	0	0	4(2.4%)
Staph aurease	2	0	0	0	2(1.2%)
Coagulase negative staphylococcus	2	0	0	0	2(1.2%)
PseudomonasAeruginosa	1	0	0	0	1(0.6%)
Total	16(9.81%)	3(1.8%)	1 (0.6%)	0	20(12.26%)

## Discussion

In our study, out of 163 patients of Sickle Cell anaemia presented with fever, many patients had localising signs of infections and respiratory system was most commonly involved with 72(44.17%) cases which include 30 cases of lobar pneumonia, 23 cases of pharyngitis, 7 cases of bronchopneumonia, 6 cases of tonsillitis, 4 cases of HRAD and 2 cases of pulmonary tuberculosis, followed by 39(23.92%) cases of gastrointestinal system, then 13(7.97%) cases of genitourinary system and 3(1.84%) cases of central nervous system(CNS). Akinyanju O, Johnson AO, 1987 in their study found, the commonest infections were pneumonia (35%), bacteraemia (32%), tonsillitis/pharyngitis (17%) and osteomyelitis (8%). (7)

Wierenga KJJ et al 2001 divided the diagnosis of fever in Sickle Cell anaemia in five groups of systemic bacterial, local bacterial, presumed viral, ACS and painful crisis and had Acute tonsillitis in 28 cases, Urinary tract infection in 4, Acute bacterial conjunctivitis in 1, Proven bacterial enteritis in 1, Upper respiratory tract infection in 27, Presumed viral enteritis in 3. 2 of asthmatic attack, Dengue haemorrhagic fever in 1, Varicella zoster infection in 1, Mumps infection in 1, Viral meningitis in 1.(6)

In present study gastrointestinal system account for 16 cases of enteric fever, 10 cases of acute gastroenteritis, 7 cases of splenic abscess, 4 cases of Hepatitis, and 2

cases of peritonitis. Dipty Jain et al in 2013, also found superficial abscess, urinary tract infection, splenic abscess and septicaemia, osteomyelitis were the causes of fever in SCD (4). Out of 16 patients of enteric fever with positive widal, salmonella typhi was isolated from 3 patients.

All 4 patients presented with features of Hepatitis were positive for Hepatitis A and out of 4 malarial patients 3 were positive for plasmodium vivax infection while 1 had plasmodium falciparum infection. Ambe JP et al 2001, stated that common problems associated with SCA patients in anaemic crisis were malaria and bacterial infections in 68 (66%) and 18 (17.3%) cases, respectively.(10)

In our study, CNS manifestations were, 2 cases of meningitis and 1 case of stroke, and CSF study in all these 3 cases were sterile. Meningitis is a major cause of death in young children with sickle cell anaemia(11). Similar results were also found in other studies also (4,6).

Urinary tract infection was found in 13(8%) patients with clinical feature and urine microscopy suggesting UTI. In 2 of these, E.coli was isolated and in 1 klebsiella pneumoniae was isolated from urine culture. Dipty et al 2013 and Wierenga KJJ et al 2001 also found UTI as cause of fever in a patients of SCD with fever(4,6)

In our study, osteomyelitis was found in 20(12.56%) patients, in which salmonella typhi was isolated from 2 patients in which 1 had concomitant enteric fever, and staphalococcus was isolated from 2 patients. Dipty Jain et al 2013 also found osteomyelitis in 21% of patients of sickle cell disease with fever(4).

Wright Jet al 1997 and Chamber JB et al 2000, found the most common offending organism in osteomyelitis was Salmonella (eight of ten cases)(12,13).

In the present study 3 patients of sickle cell anaemia with fever found to be HIV positive. Kourtis AP et al 2007 in their study, stated that hospitalized children with SCD and HIV infection have higher odds of infection than those with SCD alone(14).

7 patients had dengue fever who were diagnosed clinically and serologically. One patient had Measles, one had mumps infection, dignosed clinically and one had septicaemia in whome E. Coli was grown in blood culture. 2 patients had peritonitis out of these1 died in whome streptococcus pneumoniae was isolated from peritoneal fluid.

In our study, bacteraemia was identified in 16 (9.81%) patients. out of which Streptococcus pneumoniae was found in 3patients, Klebsiella pneumoniae was positive in 3 patients of bacteremia, E.coli was found in 2 patients of bacteremia, Salmonella typhi was found in 3patients, Staphalococcus aurease and Coagulase negative staphylococcus both of each were found in 2-2 patients of bacteremia. Pseudomonas auroginosa was found in 1 patient of bacteremia.

Williams TN et al 2009, In their study, the organisms most commonly isolated from children with sickle-cell anaemia were Streptococcus pneumoniae (44/108 isolates; 41%), non-typhi Salmonella species (19/108; 18%), Haemophilus influenzae type b (13/108; 12%), Acinetobacter species (seven of 108; 7%), and Escherichia coli (seven of 108; 7%) (15). Wierenga KJJ et al 2001, found enteric Gram negative organisms accounted for 50% of positive blood cultures: Bacteraemia was found in 10 (6.1%)patients, three Streptococcus pneumoniae, two Haemophilus influenzae type b, two Salmonella sp, one Escherichia coli, one Entero bacter sp, and one Acineto bacter sp (6). West et al 1994 and Wong WY et al 1992 also found that Streptococcus pneumoniae was the most common pathogen (26,27).

Akinyanju Oet al 1987 (7) found predominant bacteria

isolated were Klebsiella sp (38%), E. coli (23%), Staph. aureus (23%), Staph. albus(23%) and Pseudomonas spp (23%), while Okuonghae HO et al 1993 found Gram-negative bacteria were the predominant organisms (70.4%) in SCA. The commonest organisms isolated were Salmonella (25.9%), Klebsiella (25.9%) and Staphylococcus aureus (22.2) (16). In the study of Dipty Jain et al, 2013 in central India bacteraemia was proved in 20 patients of 56 acute febrile illness in which Staphylococcus aureus (n=8) and Gram negative bacteria (n=12) were responsible for all cases of bacteraemia. The isolated Gram negative bacteria included Escherichia coli (n=5), Klebsiellapneumoniae(n=5) and Pseudomonas aeruginosa (n=2) (4).

In our study, urine culture was positive in 3(1.84%) cases out of 13 UTI patients in which 2 had E.coli infection while 1 had Klebsiella infection, Wierenga KJJ et al<sup>(6)</sup> 2001, also found positive urine culture in (4/165)2.4%, which is consistent with our findings. We found 1 positive peritoneal fluid culture in which Streptococcus pneumoniae was isolated. Culture of cerebrospinal fluid(CSF) were sterile in all the 3 patients.

### Conclusion:

Most patients with homozygous sickle cell disease presenting with fever >100°F had no evidence of bacterial infection, and the fever was assumed to be attributable to viral or atypical organisms. However, these events remain clinically important because more serious pathology such as bacteraemia cannot be excluded. Despite the paucity of invasive bacterial disease in this study, it is wise to treat all febrile children with sickle cell disease with antibiotics pending the results of blood culture.

Hence we conclude that, the respiratory system is the most common one to get involved in a patients of sickle cell anaemia with fever and pneumonia is the most common diagnosis in it. Streptococcal pneumoniae along with Klebsiella pneumoniae and E. coli were found to be the common pathogens causing a significant febrile illness and bacteremia in patients of sickle cell anemia.

Acknowledgment: Data Collection done by Dr. Bhagyashree Tirpude and Dr. Shamama Subuhi. Analysis and manuscript preparation done by Dr.



Bhagyashree and all research work was done under the guidance of Dr. Dipak Madavi.

## Bibliography

- Graham R. Searjeant. Distribution of sickle cell disease. In : Graham R. Searjeant Sickle cell disease, third edition. Pg-23. Oxford university press, 2001.
- Rao VR. Genetics and epidemiology of sickle cell anemia in India. Indian J Med sci 1988; 42 : 218-22.
- Jain DL, Sarathi V, Upadhye D. Gulhane R. Nandkarni AH, Ghosh K et al. Newborn screening shows a high incidence of sickle cell anemia in Central India. Hemoglobin 2012; 36 : 316-22.
- Dipty Jain, A.S. Bagul, Maulik Shah & Vijaya Sarathil. Morbidity pattern in hospitalized under five children with sickle cell disease. Indian J Med Res 138, September 2013, pp 317-321.
- K Swarnkar, A Kale, B Lakhkar. Clinico-epidemiological and hematological profile of sickle cell anemia with special reference to penicillin prophylaxis in a rural hospital of Central India. The Int. Journal of Epidemiology. 2010 Volume 9 Number 2.
- K J JWierenga, I R Hambleton, R M Wilson, H Alexander, B E Serjeant, G R Serjeant. Significance of fever in Jamaican patients with homozygous sickle cell disease. Arch Dis Child 2001;84 : 156-159
- Akinyanju O, Johnson AO. Acute illness in Nigerian children with sickle cell anaemia. Ann Trop Paediatr. 1987 ; 7:181-186
- McIntosh S, Rooks Y, Ritchey AK, Pearson HA. Fever in children with sickle cell disease. J Pediatr. 1980 Feb;96(2):199-204.[PubMed]
- Leikin SL, Gallagher D, Kinney TR, Sloane D, Klug P, Rida 6. W. Mortality in children and adolescent with sickle cell disease. Cooperative study of sickle cell disease. Pediatrics 1989 ; 84 : 500-8
- Ambe JP, Fatunde JO, Sodeinde OO. Associated morbidities in children with sickle-cell anaemia presenting with severe anaemia in a malarious area. Trop Doct. 2001 Jan; 31(1):26-7
- McIntosh S, Rooks Y, Ritchey AK, Pearson HA. Fever in young children with sickle cell disease. J Pediatr. 1980 Feb ; 96(2) : 199-204 (PubMed).
- Wright J, Thomas P, Serjeant GR. Septicemia caused by Salmonella infection : an overlooked complication of sickle cell disease. J Pediatr. 1997 Mar ; 130(3):394-399 (PubMed).
- Chambers JB, Forsythe DA, Bertrand SL, Iwinski HJ, Stefik DE. Retrospective review of osteoarticular infections in a pediatric sickle cell age group. J PediatrOrthop. 2000;20:682-685
- Kourtis AP, Bansil P, Johnson C, Meikle SF, Posner SF, Jamieson DJ. Children with sickle cell disease and human immunodeficiency virus-1 infection use of inpatient care services in the United States. Pediatr Infect Dis J. 1997;26:406-410
- Williams TN, Uyoga S, Macharia A, Ndila C, McAuley CF, Opi DH, Mwarumba S, Makani J, Komba A, Ndiritu MN, Sharif SK, Marsh K, Berkley JA, Scott JA. Bacteraemia in Kenyan children with sickle cell anaemia: a reterospective cohort and case control study. Lancet. 2009 Oct 17;374(9698):1364-70.doi: 10.1016/S0140-6736(09)61374-X. Epub 2009 Sep 9.
- Okuonghae HO, Nwankwo MU, Offor EC. Pattern of bacteremia in febrile children with sickle cell anemia. Ann Trop Paediatr. 1993;13(1):55-64.[PubMed]
- Catherine Booth Baba Inusa Stephen K. Obaro. Infection in sickle cell disease : A review. Int J Infect Dis. 2010 Jan.
- Robinson Mg, Watson Rj. Pneumococcal Meningitis in Sickle-Cell Anemia. N Engl J Med. 1966;274(18):1006-8
- Eeckels R, Gatti F, Renoirte Am. Abnormal Distribution Of Hemoglobin Genotypes In

- Negro Children With Severe Bacterial Infections. *Nature*. 1967;216(5113):382.
20. Pearson Ha. Sickel Cell Anemia And Severe Infections Due To Encapsulated Bacteria. *J Infect Dis*. 1977;136 (Suppl):S25-30.
21. Patel AB, Athvale AM. Sickel cell disease in central India. *Indian J Pediatr* 2004; 71:789-93
22. Sadarangani M, Makani J, Komba AN, Ajala-Agbo T, Newton CR, Marsh K, et al. An observational study of children with sickel cell disease in Kilifi, Kenya. *Br J Haematol*. 2009;146(6):675-82. Epub 2009 Jul 23.
23. Lieberman L, Kirby M, Ozolins L, Mosko J, Friedman J. Initial presentation of unscreened children with sickel cell disease : the Toronto experience. *Pediatr Blood Cancer*. 2009;53(3):397-400
24. Al-Saqladi AW, Delpisheh A, Bin-Gadeem H, Brabin BJ. Clinical profile of sickel cell disease in Yemeni children. *Ann Trop Paediatr*. 2007;27(4):253-9.
25. Hawasawi ZM, Nabi G, AL Magamci MS, Awad KS. Sickel cell disease in childhood in Madina. *Ann Saudi Med*. 1998
26. West TB, West DW, Ohene-Frempong K. The presentation, frequency and outcome of bacteremia among children with sickel cell disease and fever. *Pediatr Emerg Care*. 1994 Jun ;10(3):141-143. (PubMed).
27. Wong WY, Overturf GD, Powars DR. Infection caused by *Streptococcus pneumoniae* in children with sickel cell disease : epidemiology, immunologic mechanisms, prophylaxis and vaccination. *Clin Infect Dis*. 1992;14:1124-1136.