

International Journal of Medical Science and Current Research (IJMSCR) Available online at: www.ijmscr.com Volume1, Issue 2, Page No: 174-176 July-August 2018



# Case Report: Ischemic Bilateral Opercular Syndrome (Foix-Chavany-Marie syndrome)

Dr Somnath Verma<sup>1</sup>, Dr Vaishali Manhas<sup>2</sup>, Dr Gurinder Pal Singh <sup>3</sup> <sup>1</sup>.Dr Somnath Verma, MD Medicine, Senior resident <sup>2,3</sup> Junior Resident, Department of Internal medicine, Government medical college,

Jammu (J&K), India , pin : 180001.

Corresponding Author: Somnath Verma,

MD Medicine ,Senior resident , Department of Internal medicine , Government medical college , jammu (J&K), India

Type of Publication: Original Research Paper Conflicts of Interest: Nil

#### ABSTRACT

Foix-Chavany-Marie syndrome" or "bilateral opercular syndrome" is a severe form of pseudobulbar palsy with dissociation of automatic-voluntary movements in the affected muscles. It is due to bilateral lesions of perisylvian cortex or subcortical connections and the major etiology is vascular. It is a rare neurological syndrome with only a few reported cases in the literature. Here we describe a case of Foix- Chavany-Marie syndrome due to two separate vascular events- an old infarct in the right opercular area and acute bleed in the left opercular area.

### Keywords: NIL

### **INTRODUCTION**

Opercular syndrome (OPS) is a rarely seen cortical type of pseudobulbar palsy that is characterized by paralysis of facial, lingual, pharyngeal, and masticatory voluntary muscles with preservation of autonomic, involuntary, and reflexive functions [1]. OPS, facio-labio-glosso-pharyngolaryngo- brachial paralysis, were first described by French physicians, Foix, in 1926 [2]. However, the first case was reported by Magnus (from Germany) in 1837 [3]. The lesions are usually located at the anterior part of the operculum, so it is also called anterior OPS. Herein, we report a 35-year-old male presenting with acute onset of aphonia, and facio-labio-glossopharyngolaryngeal paralysis with preservation of involuntary and emotional activities.

### 2. Case Report

A 35-year-old right-handed smoker hypertensive male presented with acute onset of sound volume loss, inability to swallow and speak, inability to move tongue, and difficulties in chewing the foods. The past medical history was unremarkable for diabetes mellitus, coronary artery disease, and benign prostate hypertrophy. His first cerebral vascular event presenting with right-sided arm and leg weakness was observed about 9 years ago and his weakness was recovered completely. However, he has not used suggested antiplatelet treatment up to now. The family history was unremarkable. In neurological assessment, he was aphonic, but his verbal and reading perception was normal. He was able to answer the questions with signs and writing. The reflexes and pupillary, corneal extraocular movements were intact. The left nasolabial sulcus disappeared. He was not able to open the mouth, protrude the tongue, show the teeth, whistle, chew, and swallow. The tongue was on the midline and immobile. No tongue atrophy, fibrillation, and deviation were observed. The taste sensation was intact. Palatal, laryngeal, blink, and gag reflexes, spontaneous smiling, and yawning were protected. The upper and lower limbs muscles strengths were

Dr. Somnath Verma et al. International Journal of Medical Science and Current Research (IJMSCR)

5/5. Deep tendon reflex was normoactive. Babinski response left side mute response and right sided flexor response . Extrapyramidal, coordination, and sensory examinations were normal. No vocal cord paralysis was detected on the laryngoscopic examination. The laboratory investigation including liver and kidney function, complete cell count, urinalysis, and erythrocyte sedimentation rate was normal. A transthoracic echocardiogram showed normal parameter. A Doppler ultrasonography of carotid and vertebral arteries and renal arteries was unremarkable.

The magnetic resonance imaging (MRI) of brain indicated first episode of left MCA terrirory infarct in 2007 (Figure 1) and NCCT head showing right MCA terriroy infarct with old left infarct (Figure 2), ischaemic stroke involving acute right temporoparietal area and insular cortex with old stroke showing gliotic changes in left temporal region bilateral chronic inferior frontal gyrus infarct (Figure 3). He was treated with 300mg acetylsalicylic acid daily and was placed nasogastric tube for feeding due to lack of swallowing function. At the end of the 3rd week, he had nonsense sounds and was able to open the mouth. However, there was no improvement in speech, chewing, and swallowing functions.



Fig 1 . old MRI Brain showing left MCA territory infarct







Fig.3; Magnetic resonance imaging sequences demonstrate right temporoparietal area and insular cortex with old stroke showing gliotic changes in left temporal region bilateral chronic inferior frontal gyrus infarct (R: Right, L: Left).

## 3. Discussion

OPS are characterized by anarthria and paralysis of the facial, pharyngeal, masticatory, tongue, laryngeal, and brachial voluntary muscles. Operculum refers to the brain cortex covering the insula, inferior frontal, pre- and postcentral, supramarginal, angular (inferior parietal), and superior temporal gyrus. OPS are particularly originated from the damages that are located in posterior part of the inferior frontal gyrus and inferior part of precentral gyrus [4]. There are several connections between bilateral precentral gyrus and cranial nerves [5, 6, 9, 10, and 12,] and

#### 

bilateral damage of these corticobulbar tracts can cause OPS.

The voluntary muscle controls of the face, tongue, and pharynx are provided by primary motor cortex and pyramidal tract, whereas the spontaneous and emotional controls are provided by thalamus, hypothalamus, and Extrapyramidal tract [5].The selective paralysis of voluntary muscle weakness in OPS is named as "autonomic-voluntary dissociation."

The etiologic factors of OPS include thrombotic or embolic multiple strokes, head trauma, tumor, developmental perisylvian dysplasia, multiple sclerosis, acute disseminated encephalomyelitis, moyamoya disease, vasculitis, and neurodegenerative disease [6]. Ischemic stroke related OPS is the most common classic type. In this type of OPS, as seen in our case, patients usually have history of one or more contralateral stroke. Unilateral lesion rarely can cause OPS, but it was postulated that those cases might have subtle contralateral lesions. The SPECT imaging study on the patients with unilateral OPS showed cerebral diaschisis that supported the hypothesis [7]. We think that combination of both old left inferior frontal gyrus lesion and the new acute infarct of the operculum contributed to the clinical picture of OPS.

Pseudobulbar paralysis in OPS is clinically distinguished from bulbar paralysis, disorders of the cranial nerves and neuromuscular junction (e.g., botulism and myasthenia gravis) by normal eye movements, preserved or hyperactive brainstem reflexes (e.g., jaw jerk), the dissociation of automatic and volitional movements of the bulbar muscles with preservation of automatic movements, and the absence of atrophy and fasciculations of the lower motor neuron innervated muscles [8].

The treatment and prognosis of OPB is related to underlying etiological factors. However, clinical improvement is usually poor. Chewing, swallowing, and speech functions do not usually recover completely. Patients with OPS have a significant risk for aspiration pneumonia. Therefore, during acute treatment and rehabilitation process, speech dysfunction and feeding are two most important issues. In our case, no improvement in speech, swallowing, and chewing functions was observed. He could produce nonsense sounds and open the mouth at the end of the 3rd weeks. Preserving patients from OPS can be possible via preventive treatment of repetitive stroke.

### **Conflict of Interests**

The authors declare that they have no conflict of interests.

### **Authors' Contribution**

The final paper has been seen and approved by all authors. The authors accept full responsibility for the design and conduct of the study, had access to the data, and controlled the decision to publish.

### Disclosure

The authors also report the absence of any significant financial support in any organization. The paper had not been published elsewhere previously.

### References

. . . . . . . . . . . . . . .

[1] M. Y. Ann, O. K. Liu, and Y. L. Wu, "Foix-Chavany-Marie syndrome," *Chinese Medical Journal*, vol. 64, no. 9, pp. 540–544, 2001.

[2] C. Foix, J. A. Chavany, and J. Marie, "Diplegie faciolinguomasticatrice d'origine cortico-sous-cortical sans paralysie des members," *Revue Neurologique*, vol. 33, pp. 214–219, 1926.

[3] A. Magnus, "Fall von aufhebung des willenseinflusses auf einige hirnnerven," *Archiv f ur Anatomie, Physiologie und Wissenschaftliche Medicin*, pp. 258–266, 1837.

[4] M. Weller, "Anterior opercular cortex lesions cause dissociated lower cranial nerve palsies and anarthria but no aphasia: foixchavany- marie syndrome and "automatic voluntary dissociation" revisited," *Journal ofNeurology*, vol. 240, pp. 199–208, 1993.

[5] C. C.Mao, B.M. Coull, L. A. C. Golper, and M. T. Rau, "Anterior operculum syndrome," *Neurology*, vol. 39, no. 9, pp. 1169–1172, 1989.

[6] M. Bakar, H. S. Kirshner, and F. Niaz, "The opercularsubopercular syndrome: four cases with review of the literature," *Behavioural Neurology*, vol. 11, no. 2, pp. 97–103, 1998.

[7] E. Kutluay, Z. Colakoglu, A. Dirlik, and K. Kumral, "Brain SPECT in anterior opercular syndrome due to a unilateral lesion," *Journal of Neurology*, vol. 243, no. 5, pp. 427–429, 1996.

[8] D. J. Lanska, "Foix-chavany-marie syndrome," MedLink Neurology, http://www.Medlink.com/medlinkcontent.asp.

Volume 1, Issue 2; July-August 2018; Page No. 174-176 © 2018 IJMSCR. All Rights Reserved