

Correction of Smile and Profile in a Patient with Bimaxillary Dentoalveolar Protrusion Using Pre-Adjusted Edgewise Mechanotherapy – A Case Report

Dr. Bhushan Jawale¹, Dr. Lishoy Rodrigues², Dr. Vijay Naik³, Dr. Veerendra Kerudi⁴, Dr. Pushkar Gawande⁵, Dr. Amit Nehete⁶

¹Professor, ²PG Student, ^{3,4}Professor and HOD, ⁵Reader, ⁶Professor
 Dept of Orthodontics and Dentofacial Orthopedics,

^{1,2,5}Sinhgad Dental College and Hospital, Vadgaon Bk, Pune, Maharashtra, India, ³Maratha Mandal Dental College and Hospital, Belgaum, Karnataka, India, ⁴ACPM Dental College and Hospital, Dhule, Maharashtra, India. ⁶MGV Dental College and Hospital, Nashik, Maharashtra, India

***Corresponding Author:**

Dr. Lishoy Rodrigues

PG Student, Dept of Orthodontics and Dentofacial Orthopedics, Sinhgad Dental College and Hospital, Vadgaon Bk, Pune, Maharashtra, India

Type of Publication: Case Report

Conflicts of Interest: Nil

ABSTRACT

This case report evaluates the management of bimaxillary dentoalveolar protrusion in a female patient having a Class I malocclusion with Pre-adjusted Edgewise bracket system. The case required extraction of 1st premolars for correction of the proclined and forwardly placed upper and lower front teeth. Clinical and cephalometric evaluation revealed skeletal Class I pattern with anterior proclination and a convex facial profile, an average to vertical growth pattern, incompetent lips, a posterior divergent face, an average overjet and overbite. Following fixed orthodontic treatment by removal of 1st premolars in the upper and lower arch with retraction of anterior segment, a marked improvement in patient's smile, facial profile and occlusion was achieved and there was a remarkable increase in the patient's confidence and quality of life. The profile changes and treatment results were demonstrated with proper case selection and good patient cooperation with fixed appliance therapy.

Keywords: Correction of Smile, profile, Pre-adjusted Edgewise bracket system, Bimaxillary dentoalveolar protrusion, Fixed Appliance Therapy, Class I malocclusion, Crowded dentition, Leptoprosopic facial form, Long face, Aesthetic Improvement, 1st Premolar Extraction, Orthodontic Camouflage, Therapeutic Extractions

INTRODUCTION

Orthodontic treatment can significantly alter and improve facial appearance in addition to correcting irregularity of the teeth. Class I malocclusion is the 2nd most prevalent malocclusion after Class II.^[1,4,5] Over the last few decades, there has been an increase in the awareness about orthodontic treatment which has led to more and more adults demanding high quality treatment in the shortest possible time with increased efficiency and reduced costs.^[2,3,7] There are many ways to treat Class I malocclusions, according to the characteristics

associated with the problem, such as antero-posterior discrepancy, age, and patient compliance.^[6] The indications for extractions in orthodontic practice have historically been controversial.^[8] On the other hand, correction of Class I malocclusions in growing patients, with subsequent dental camouflage to mask the skeletal discrepancy, can involve either retraction by non-extraction means simply by utilizing the available spaces or by extractions of premolars.^[9,10] Lack of crowding or cephalometric discrepancy in the mandibular arch is an indication of

2 premolar extraction. Fortunately, in some instances satisfactory results with an exceptional degree of correction can be achieved without extraction of permanent premolars ^[11-13]. This case presents the correction of a Bimaxillary dentoalveolar protrusion with a Class I malocclusion in an adult female patient with severely proclined maxillary and mandibular anterior teeth merely simply by executing extraction of maxillary and mandibular 1st premolars followed by fixed appliance therapy using Pre-adjusted Edgewise bracket system. The Extraction protocol shown in this case is indicative of how a convex unesthetic facial profile can be converted into an Orthognathic pleasant profile by routine fixed Orthodontic treatment with extraction of 4 premolars followed by retraction and closure of spaces.

CASE REPORT

PRE TREATMENT EXTRA ORAL PHOTOGRAPHS



EXTRA-ORAL EXAMINATION


A 26 year old female patient presented with the chief complaint of forwardly placed upper and lower front teeth and excessive show of front teeth. On Extraoral examination, the patient had a convex facial profile, grossly symmetrical face on both sides with slightly retruded chin, incompetent lips ,shallow mentolabial sulcus and an acute Nasolabial Angle , a Leptoprosopic facial form, Dolicocephalic head form, average width of nose and mouth, a consonant smile arc and posterior divergence of face along with an average to vertical growth pattern. The patient had no relevant prenatal, natal, postnatal history, history of habits or a family history. On Smiling, there was excessive show of maxillary anterior teeth with slightly broad buccal corridor spaces. The patient had a toothy smile and an unaesthetic facial profile. The patient was very dissatisfied with her smile.



INTRA-ORAL EXAMINATION

Intraoral examination on frontal view shows presence of an average overjet and overbite with lower dental midline shifted to the right by 0.5mm. On lateral view the patient shows the presence of Class I incisor relationship and a Class I canine and molar relationship bilaterally. Patient has proclined and forwardly placed upper and lower anterior teeth.

PRE TREATMENT INTRA ORAL PHOTOGRAPHS





PRE TREATMENT CEPHALOMETRIC READINGS

PARAMETERS	PRE- TREATMENT
SNA	82°
SNB	80°
ANB	2°
WITS	1mm(AO ahead of BO)
MAX. LENGTH	100mm
MAN. LENGTH	96mm
IMPA	113°
NASOLABIAL ANGLE	87°
U1 TO NA DEGREES	43°
U1 TO NA mm	6mm
L1 TO NB DEGREES	36°
L1 TO NB mm	6mm
U1/L1 ANGLE	106°
FMA	28°
Y AXIS	75°

DIAGNOSIS

This 26 year old female patient was diagnosed with a Class I malocclusion on a Class I skeletal base , an average to vertical growth pattern, average overjet and overbite with a lower midline shift to the right by 0.5mm, proclined upper and lower incisors, protruded upper and lower lips, a slightly retruded chin, moderately deep mentolabial sulcus, increased lip strain, incompetent lips with an acute Nasolabial angle and a convex facial profile with a posteriorly divergent face.

LIST OF PROBLEMS

1. Proclined maxillary and mandibular anterior teeth
2. Convex facial profile
3. Decreased Nasolabial angle
4. Incompetent lips
5. Increased lip strain
6. Non coincident dental midlines

TREATMENT OBJECTIVES

1. To correct proclined maxillary and mandibular anterior teeth
2. To correct the posterior divergence of face
3. To correct the decreased Nasolabial angle
4. To correct the dental midlines
5. To decrease the lip strain
6. To achieve a pleasing smile and a pleasing profile

TREATMENT PLAN

- Extraction of 14, 24, 34 and 44
- Fixed appliance therapy with Pre-adjusted Edgewise bracket system
- Initial leveling and alignment with 0.012", 0.014", 0.016", 0.018", 0.020" NiTi arch wires following sequence A of MBT

- Retraction and closure of spaces by use of 0.019" x 0.025" rectangular NiTi followed by 0.019" x 0.025" rectangular stainless steel wires.
- Final finishing and detailing with 0.014" round stainless steel wires
- Retention by means of Begg's Wrap-around retainers along with lingual bonded retainers in the upper and lower arch

TREATMENT PROGRESS

Complete bonding & banding in both maxillary and mandibular arch was done, using Pre-adjusted Edgewise bracket system. Initially a 0.012" NiTi wire was used which was followed by 0.014", 0.016", 0.018", 0.020" NiTi arch wires following sequence A of MBT. After 6 months of alignment and leveling NiTi round wires were discontinued. Retraction and closure of spaces was then started by use of 0.019" x 0.025" rectangular NiTi followed by 0.019" x 0.025" rectangular stainless steel wires. Reverse curve of spee in the lower arch and exaggerated curve of spee in the upper arch was incorporated in the heavy archwires to prevent the excessive bite deepening during retraction process and also to maintain the normal overjet and overbite. Group A anchorage was maintained in the upper and lower arch. Retraction and closure of spaces was done with the help of Elastomeric chains delivering light continuous forces and replaced after every 4 weeks due to force decay and reduction in its activity. Finally light settling elastics were given with rectangular steel wires in lower arch and 0.012" light NiTi wire in upper arch for settling , finishing, detailing and proper intercuspation. The increased overjet was corrected with an ideal occlusion at the end of the fixed appliance therapy. Also the profile of the patient improved significantly to more Orthognathic with a pleasant and consonant smile arc on smiling. Also, the Nasolabial angle improved significantly at the end of treatment.

POST TREATMENT CEPHALOMETRIC READINGS

PARAMETERS	POST-TREATMENT
SNA	81°
SNB	79°
ANB	2°
WITS	1mm(AO ahead of BO)
MAX. LENGTH	99mm
MAN. LENGTH	95mm
IMPA	93°
NASOLABIAL ANGLE	107°
U1 TO NA DEGREES	27°
U1 TO NA mm	2mm
L1 TO NB DEGREES	26°
L1 TO NB mm	2mm
U1/L1 ANGLE	132°
FMA	27°
Y AXIS	74°

POST TREATMENT EXTRA ORAL PHOTOGRAPHS





POST TREATMENT INTRA ORAL PHOTOGRAPHS



DISCUSSION

A well-chosen individualized treatment plan, undertaken with sound biomechanical principles and appropriate control of orthodontic mechanics to execute the plan is the surest way to achieve predictable results with minimal side effects. Treatment of a Class I malocclusion with extractions of all 1st premolars is challenging. Class I malocclusion with Bimaxillary Dentoalveolar protrusion might have any number of a combination of the skeletal and dental components. Hence, identifying and understanding the etiology and expression of Class I malocclusion and identifying differential diagnosis is helpful for its correction. The patient's chief complaint was forwardly placed upper and lower front teeth with excessive show of front teeth. The case was of a clear bimaxillary dentoalveolar protrusion with severely proclined upper and lower anterior dentition. The selection of orthodontic fixed appliances is dependent upon several factors which can be categorized into patient factors, such as age and compliance, and clinical factors, such as preference/familiarity and laboratory facilities. In this case we choose to use Pre-adjusted Edgewise bracket system. The execution of all 1st premolar extraction followed by Fixed appliance therapy appropriately resulted in an improvement in the patient's convex profile in this case. The most important point to be highlighted here is the decision to extract the premolars. After analyzing the case

thoroughly and reading all pretreatment cephalometric parameters along with evaluating the patients profile clinically, a decision was made of extracting the 1st premolars. Proximal stripping with retraction and closure of spaces could not be executed in this case as this would not address all the patient problems at the end of the treatment. The patient had excessive proclination of maxillary and mandibular anterior teeth. Also the patient had a convex profile with an acute nasolabial angle and a severely decreased Inter-incisal angle. All these findings made it essentially imperative to extract all 1st premolars. This case could not be managed by non-extraction or proximal stripping. Extractions also very efficiently improved the patient's profile, changing it from being slightly convex to more orthognathic at the end of the treatment. There was improvement in occlusion, smile arc, profile and position of chin. Successful results were obtained after the fixed Pre-adjusted Edgewise appliance therapy within a stipulated period of time. The overall treatment time was 17 months. After this active treatment phase, the profile of this 26 year old female patient improved significantly as seen in the post treatment Extra-oral photographs. Removable Begg's retainers were then delivered to the patient along with fixed lingual bonded retainers in upper and lower arch. One year follow up records were taken and did not reveal any drastic untoward changes in the patients smile and profile.

POST TREATMENT CEPHALOMETRIC READINGS

PARAMETERS	POST-TREATMENT
SNA	81°
SNB	79°
ANB	2°
WITS	1mm(AO ahead of BO)
MAX. LENGTH	99mm
MAN. LENGTH	95mm
IMPA	94°
NASOLABIAL ANGLE	106°
U1 TO NA DEGREES	28°
U1 TO NA mm	2mm

L1 TO NB DEGREES	26°
L1 TO NB mm	2mm
U1/L1 ANGLE	130°
FMA	27°
Y AXIS	73°

1 YEAR FOLLOW UP EXTRA ORAL PHOTOGRAPHS




COMPARISON OF PRE AND POST TREATMENT AND ONE YEAR FOLLOW-UP CEPHALOMETRIC READINGS

PARAMETERS	PRE- TREATMENT	POST-TREATMENT	ONE YEAR FOLLOW - UP
SNA	82°	81°	81°
SNB	80°	79°	79°
ANB	2°	2°	2°
WITS	1mm(AO ahead of BO)	1mm(AO ahead of BO)	1mm(AO ahead of BO)
MAX. LENGTH	100mm	99mm	99mm
MAN. LENGTH	96mm	95mm	95mm
IMPA	113°	93°	94°
NASOLABIAL ANGLE	87°	107°	106°
U1 TO NA DEGREES	43°	27°	28°
U1 TO NA mm	6mm	2mm	2mm

L1 TO NB DEGREES	36°	26°	26°
L1 TO NB mm	6mm	2mm	2mm
U1/L1 ANGLE	106°	132°	130°
FMA	28°	27°	27°
Y AXIS	75°	74°	73°

PROFILE CHANGES PRE AND POST TREATMENT AND AFTER 1 YEAR



CONCLUSION

This case report shows how Bimaxillary Dentoalveolar Protrusion case can be managed with Extraction of 4 premolars by means of appropriate use of Pre-adjusted Edgewise bracket system and efficient conservation of anchorage at the same time. The planned goals set in the pretreatment plan were successfully attained. Treatment of the proclined and forwardly placed upper and lower anterior teeth included the retraction and retroclination of maxillary and mandibular incisors with a resultant decrease in soft tissue procumbency and facial convexity. The profile changed from slightly convex to orthognathic. The maxillary and mandibular teeth were found to be esthetically satisfactory in the line of occlusion. Patient had an improved smile and profile. The correction of the malocclusion was achieved, with a significant improvement in the patient aesthetics and self-esteem. The patient was very satisfied with the result of the treatment.

REFERENCES

1. Bhushan Jawale D, Rodrigues L, Keluskar KM, Jatti R, Belludi A, Hattarki R. Treatment of a growing male having a recessive mandible with removable myofunctional appliance therapy followed by fixed orthodontic treatment
2. Jawale B, Rodrigues L, Garde JB, Belludi A, Patil A, Palande P. Interdisciplinary collaboration of orthodontics and oral and maxillofacial surgery for the correction of severe class III skeletal pattern in an adult male with an hapsburg jaw-A case report on surgical orthodontics. IP Indian Journal of Orthodontics and Dentofacial Research. 2020 Sep 15;6(3):149-56.
3. Lishoy R, Priyal R, Jamenis SC, Jawale B, Mahajan N. A survey to assess the knowledge and attitude of adults from the age group of

- 18 to 35 Years towards comprehensive orthodontic treatment-A questionnaire based study on adult orthodontics. IP Indian Journal of Orthodontics and Dentofacial Research. 2020 Nov 15;6(4):255-63.
4. Bhushan Jawale D, Rodrigues L, Naik V, Kerudi V, Chaudhary A, Nehete A. Management of a non growing adult borderline extraction case of a patient having a Class II Division 1 malocclusion by non extraction protocol for aesthetic improvement: A case report on adult orthodontics.
 5. Jawale B, Lishoy R, Belludi A, Pharande A, Hattarki R, Prasad L. Correction of bimaxillary dentoalveolar protrusion in a growing male with class I malocclusion by extraction of premolars and profile improvement using conventional fixed orthodontic treatment-A case report on orthodontic camouflage. IP Indian Journal of Orthodontics and Dentofacial Research. 2020 Sep 15;6(3):157-62.
 6. Rodrigues L, Jawale B, Kadam A, Rajani P. Single phase correction of tongue thrust habit alongside fixed orthodontic treatment for closure of spaced dentition and midline diastema in a male patient with class I malocclusion without need for a two phase appliance therapy-A case report. IP Indian Journal of Orthodontics and Dentofacial Research. 2020 Sep 15;6(3):163-9.
 7. Rodrigues L, Jawale B, Jamenis S, Sadhunavar T. Application of magnets in orthodontics-A review. IP Journal of Surgery and Allied Sciences. 2020 Nov 15;2(3):50-7.
 8. Rodrigues L, Jamenis SC, Jawale B, Patil R, Sadhunavar T. An assessment of knowledge and application of lingual orthodontics among orthodontists in their routine clinical practice. IP Journal of Surgery and Allied Sciences. 2020 Nov 15;2(3):89-94.
 9. Rodrigues L, Jamenis SC, Jawale B, Patil S, Garcha V. A questionnaire study to assess and evaluate the common gingival problems faced by patients undergoing fixed orthodontic treatment. IP International Journal of Maxillofacial Imaging. 2021 Jan 15;6(4):101-7.
 10. Jawale B, Rodrigues L, Shinde K, Kangane S, Hattarki R, Mhatre S. Rhinoplasty, septoplasty and genioplasty with fixed orthodontic mechanotherapy for non-surgical correction of a patient with “Long face syndrome” Having a class III malocclusion on a class II skeletal jaw base-A case report. IP Indian Journal of Orthodontics and Dentofacial Research. 2020 Sep 15;6(3):170-6.
 11. Jawale B, Rodrigues L, Keluskar KM, Patil S, Belludi A, Patil A. Forsus fixed functional appliance therapy for dentoalveolar and profile correction-A case report. IP Indian Journal of Orthodontics and Dentofacial Research. 2020 Nov 15;6(4):264-70.
 12. Rashi L, Priyal R, Marisca P, Aljeeta K. An assessment of common concerns of 2nd year post graduate students pursuing MDS In orthodontics and dentofacial orthopedics, due to the COVID-19 lockdown.
 13. Rodrigues L, Jawale B, Kaluskar A, Jadhav B, Kadam A, Shaikh A, Gurav K, Borchate T. Molar Banding or Bonding? What do Orthodontists Prefer in Routine Clinical Practice?